arXiv:1704.03586v1 [math.CA] 12 Apr 2017

BILINEAR SPHERICAL MAXIMAL FUNCTION

J. A. BARRIONEVO, LOUKAS GRAFAKOS, DANQING HE, PETR HONZÍK, AND LUCAS OLIVEIRA

ABSTRACT. We obtain boundedness for the bilinear spherical maximal function in a range of exponents that includes the Banach triangle and a range of L^p with $p<1$. We also obtain counterexamples that are asymptotically optimal with our positive results on certain indices as the dimension tends to infinity.

1. INTRODUCTION

Let σ be surface measure on the unit sphere. The spherical maximal function

$$\mathscr{M}(f)(x)=\sup_{t>0}\Big|\int_{|y|=1} f(x-ty)d\sigma(y)\Big|, \tag{1}$$

was first studied by Stein [19] who provided a counterexample showing that it is unbounded on $L^p(\mathbb{R}^n)$ for $p\le\frac{n}{n-1}$ and obtained the a priori inequality $\|\mathscr{M}(f)\|_{L^p(\mathbb{R}^n)}\le C_{p,n}\|f\|_{L^p(\mathbb{R}^n)}$ when $n\ge 3$, $p\in(\frac{n}{n-1},\infty)$ for smooth functions f; see also the account in [20, Chapter XI]. The extension of this result to the case $n=2$ was established about a decade later by Bourgain [1].

In addition to Stein and Bourgain, other authors have studied the spherical maximal function; for instance see [5], [3], [17], [16], and [18]. Among the techniques used in these works, we highlight that of Rubio de Francia [17], in which the L^p boundedness of (1) is reduced to certain L^2 estimates obtained by Plancherel's theorem. Extensions of the spherical maximal function to different settings have also been established by several authors: for instance see [4], [2] [12], [7] and [15].

In this work we study the bi(sub)linear spherical maximal function defined in (2), which was introduced and first studied by [8]. In the bilinear setting the role of the crucial $L^2\to L^2$ estimate is played by an $L^2\times L^2\to L^1$, and obviously Plancherel's identity cannot be used on L^1. We overcome the lack of orthogonality on L^1 via a wavelet technique introduced by three of the authors in [10] in the study of certain bilinear operators; on this approach

The fourth author was supported by the ERC CZ grant LL1203 of the Czech Ministry of Education. The second author acknowledges the support of Simons Foundation and of the University of Missouri Research Board and Research Council.

see [11], [14]. It is worth mentioning a related interesting recent paper [13], where the authors studied the bilinear circular average when $n = 1$. Our object of study here is the bi(sub)linear spherical maximal function

$$\mathcal{M}(f,g)(x) = \sup_{t>0} \Big| \int_{\mathbb{S}^{2n-1}} f(x-ty)g(x-tz)d\sigma(y,z) \Big| \tag{2}$$

initially defined for Schwartz functions f, g on $\mathbb{R}^n$. Here σ is surface measure on the $2n-1$-dimensional sphere. We are concerned with bounds for $\mathcal{M}$ from a product of Lebesgue spaces $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to another Lebesgue space $L^p(\mathbb{R}^n)$, where $1/p = 1/p_1 + 1/p_2$. The main result of this article is the following:

Theorem 1. *Let $n \geq 8$ and let $\delta_n = (2n-15)/10$. Then the bilinear maximal operator $\mathcal{M}$, when restricted to Schwartz functions, is bounded from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$ with $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$ for all indices $(\frac{1}{p_1}, \frac{1}{p_2}, \frac{1}{p})$ in the open rhombus with vertices the points $\vec{P}_0 = (\frac{1}{\infty}, \frac{1}{\infty}, \frac{1}{\infty})$, $\vec{P}_1 = (1, \frac{1}{\infty}, 1)$, $\vec{P}_2 = (\frac{1}{\infty}, 1, 1)$ and $\vec{P}_3 = (\frac{1+2\delta_n}{2+2\delta_n}, \frac{1+2\delta_n}{2+2\delta_n}, \frac{1+2\delta_n}{1+\delta_n})$.*

Once Theorem 1 is known, it follows that $\mathcal{M}$ admits a bounded extension from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$ for indices in the open rhombus of Theorem 1 (for such indices we have $p_1, p_2 < \infty$). Indeed, given $\{f_j\}_j$ Schwartz functions converging to f in L^{p_1} and $\{g_k\}_k$ Schwartz functions converging to g in L^{p_2}, we have that

$$\|\mathcal{M}(f_j, g_j) - \mathcal{M}(f_{j'}, g_{j'})\|_{L^p} \leq \big\|\mathcal{M}(f_j - f_{j'}, g_j) + \mathcal{M}(f_j, g_j - g_{j'})\big\|_{L^p}.$$

It follows from this that the sequence $\{\mathcal{M}(f_j, g_j)\}_j$ is Cauchy in $L^p(\mathbb{R}^n)$ and hence it converges to a value which we also call $\mathcal{M}(f,g)$. This is the bounded extension of $\mathcal{M}$ from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$. In order to pass to the maximal function defined on $L^{p_1} \times L^{p_2}$, it is also possible to used the technique desribed in [20, page 508].

Concerning dimensions smaller than 8, we have positive answers in the Banach range in next section.

2. The Banach range in dimensions $n \geq 2$

Proposition 2. *Let $n \geq 2$. Then $\mathcal{M}$ maps $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$ when $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$, $1 < p_1, p_2 \leq \infty$, and $1 < p \leq \infty$.*

Proof. We show that $\mathcal{M}$ is bounded on the intervals $[\vec{P}_0, \vec{P}_1)$ and $[\vec{P}_0, \vec{P}_2)$, where $\vec{P}_1$ and $\vec{P}_2$ are as in Theorem 1. Then the claimed assertion follows by interpolation. If one function, for instance the second one g, lies in L^∞, matters reduce to the $L^p(\mathbb{R}^n)$ boundedness of the maximal operator

$$\mathcal{M}^0(f)(x) = \sup_{t>0} \int_{\mathbb{S}^{2n-1}} |f(x-ty)| d\sigma(y,z),$$

since $\mathcal{M}(f,g)(x) \le \|g\|_{L^\infty}\mathcal{M}^0(f)(x)$. This expression inside the supremum is a Fourier multiplier operator of the form

$$\int_{\mathbb{R}^{2n}} \widehat{|f|}(\xi)\delta_0(\eta)\widehat{d\sigma}(t\xi,t\eta)e^{2\pi ix\cdot(\xi+\eta)}d\xi d\eta = \int_{\mathbb{R}^n} \widehat{|f|}(\xi)\widehat{d\sigma}(t\xi,0)e^{2\pi ix\cdot\xi}d\xi$$

where δ_0 is the Dirac mass and

$$\widehat{d\sigma}(t(\xi,0)) = 2\pi\,\frac{J_{n-1}(2\pi t|(\xi,0)|)}{|t(\xi,0)|^{n-1}}.$$

The multiplier $\widehat{d\sigma}(\xi,0)$ is smooth everywhere and decays like $|\xi|^{-(n-\frac{1}{2})}$ as $|\xi|\to\infty$ and its gradient has a similar decay.

The following result is in [17, Theorem B] (see also [6]):

Theorem A. *Let $m(\xi)$ be a $\mathcal{C}^{[n/2]+1}(\mathbb{R}^n)$ function that satisfies $|\partial^\gamma m(\xi)| \le (1+|\xi|)^{-a}$ for all $|\gamma| \le [n/2]+1$ with $a \ge (n+1)/2$. Then the maximal operator*

$$f \mapsto \sup_{t>0} \left|\left(\widehat{f}(\xi)m(t\xi)\right)^\vee\right|$$

maps $L^p(\mathbb{R}^n)$ to itself for $1<p<\infty$.

In order to have $n-\frac{1}{2} \ge \frac{n+1}{2}$ we must assume that $n \ge 2$. It follows from Theorem A that $\mathcal{M}^0$ is bounded on L^p when $1<p\le\infty$ and $n\ge 2$. This completes the proof of Proposition 2.

□

3. The point $(2,2,1)$

Next we turn to the main estimate of this article which concerns the point $L^2\times L^2 \to L^1$, i.e., the estimate $\|\mathcal{M}(f,g)\|_{L^1} \le \|f\|_{L^2}\|g\|_{L^2}$.

Proposition 3. *If ψ is in $\mathcal{C}_0^\infty(\mathbb{R}^{2n})$, then the maximal function*

$$M(f,g)(x) = \sup_{t>0}\left|\int_{\mathbb{R}^{2n}} \widehat{f}(\xi)\widehat{g}(\eta)\psi(t\xi,t\eta)e^{2\pi ix\cdot(\xi+\eta)}d\xi d\eta\right|$$

satisfies that for any $1<p_1,p_2<\infty$ and $1/p = 1/p_1+1/p_2$, there exists a constant C independent f and g such that

$$\|M(f,g)\|_{L^p(\mathbb{R}^n)} \le C\|f\|_{L^{p_1}(\mathbb{R}^n)}\|g\|_{L^{p_2}(\mathbb{R}^n)}$$

The proof of Proposition 3 is standard and is omitted. Next, we decompose $\mathcal{M}$. We fix $\varphi_0 \in \mathcal{C}_0^\infty(\mathbb{R}^{2n})$ such that $\chi_{B(0,1)} \le \varphi_0 \le \chi_{B(0,2)}$ and we let $\varphi(\xi,\eta) = \varphi_0((\xi,\eta)) - \varphi_0(2(\xi,\eta))$. For $j\ge 1$ define

$$m_j(\xi,\eta) = \widehat{d\sigma}(\xi,\eta)\varphi(2^{-j}(\xi,\eta))$$

and for $j=0$ define $m_0(\xi,\eta)=\widehat{d\sigma}(\xi,\eta)\varphi_0(\xi,\eta)$. Then we have

$$\widehat{d\sigma}=m=\sum_{j\geq 0} m_j$$

where $\widehat{d\sigma}(\xi,\eta)=2\pi\frac{J_{n-1}(2\pi(\xi,\eta))}{|(\xi,\eta)|^{n-1}}$. Setting

$$\mathcal{M}_j(f,g)(x)=\sup_{t>0}\left|\int_{\mathbb{R}^{2n}}\widehat{f}(\xi)\widehat{g}(\eta)m_j(t\xi,t\eta)e^{2\pi ix\cdot(\xi+\eta)}d\xi d\eta\right|,$$

we have the pointwise estimate

$$\mathcal{M}(f,g)(x)\leq\sum_{j\geq 0}\mathcal{M}_j(f,g)(x),\qquad x\in\mathbb{R}^n. \tag{3}$$

Proposition 4. *For $n\geq 8$, there exist positive constants C and $\delta_n=\frac{n}{5}-\frac{3}{2}$ such that for all $j\geq 1$ and all functions $f,g\in L^2(\mathbb{R}^n)$ we have*

$$\|\mathcal{M}_j(f,g)\|_{L^1}\leq C\,j\,2^{-\delta_n j}\|f\|_{L^2}\|g\|_{L^2}. \tag{4}$$

Proposition 4 will be proved in the next section. In the remaining of this section we state and prove a lemma needed for its proof.

Lemma 5. *Suppose that $\sigma_1(\xi,\eta)$ is defined on $\mathbb{R}^{2n}$ and for some $\delta>0$ it satisfies:*

(i) for any multiindex $|\alpha|\leq M=4n$, there exists a positive constant C_α independent of j such that $\|\partial^\alpha(\sigma_1(\xi,\eta))\|_{L^\infty}\leq C_\alpha 2^{-j\delta}$,

(ii) supp $\sigma_1\subset\{(\xi,\eta)\in\mathbb{R}^{2n}:|(\xi,\eta)|\sim 2^j, c_1 2^{-j}\leq\frac{|\xi|}{|\eta|}\leq c_2 2^j\}$.

Then $T(f,g)(x):=\int_0^\infty|T_{\sigma_t}(f,g)(x)|\frac{dt}{t}$ is bounded from $L^2(\mathbb{R}^n)\times L^2(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$ with bound at most a multiple of $j\|\sigma_1\|_{L^2}^{4/5}2^{-j\delta/5}$, where $\sigma_t(\xi,\eta)=\sigma_1(t\xi,t\eta)$.

Proof of Lemma 5. A crucial tool in the proof of Lemma 5 is the following result [10, Corollary 8]:

Proposition B. *Let $m\in L^2(\mathbb{R}^{2n})$ and $C_M>0$ satisfy $\|\partial^\alpha m\|_{L^\infty}\leq C_M$ for each multiindex $|\alpha|\leq M=16n$. Then the bilinear operator T_m associated with the multiplier m satisfies*

$$\|T_m\|_{L^2\times L^2\to L^1}\leq C C_M^{1/5}\|m\|_{L^2}^{4/5}.$$

Using Proposition B, setting $\widehat{f^j}=\widehat{f}\chi_{\{c_1\leq|\xi|\leq c_2 2^{j+1}\}}$, by the support of σ_1 we obtain that

$$\|T_{\sigma_1}(f,g)\|_{L^1}\leq C\|\sigma_1\|_{L^2}^{4/5}2^{-j\delta/5}\|f^j\|_{L^2}\|g^j\|_{L^2}.$$

Notice that $T_{\sigma_t}(f,g)(x) = t^{-2n} T_{\sigma_1}(f_t, g_t)(\frac{x}{t})$, where $\widehat{f_t}(\xi) = \widehat{f}(\xi/t)$. Then

$$\begin{aligned}\|T_{\sigma_t}(f,g)\|_{L^1} &\leq C\|\sigma_1\|_{L^2}^{4/5} 2^{-j\delta/5} t^{-n} \|\widehat{f}(\xi/t)\chi_{E_{j,0}}\|_{L^2} \|\widehat{g}(\eta/t)\chi_{E_{j,0}}\|_{L^2} \\ &= C\|\sigma_1\|_{L^2}^{4/5} 2^{-j\delta/5} \|\widehat{f}\chi_{E_{j,t}}\|_{L^2} \|\widehat{g}\chi_{E_{j,t}}\|_{L^2},\end{aligned}$$

where $E_{j,t} = \{\xi \in \mathbb{R}^n : \frac{c_1}{t} \leq |\xi| \leq \frac{2^j c_2}{t}\}$.

As a result we obtain

$$\begin{aligned}&\int_{\mathbb{R}^n} \int_0^\infty |T_{\sigma_t}(f,g)| \frac{dt}{t} dx \\ \leq & C\|\sigma_1\|_{L^2}^{4/5} 2^{-j\delta/5} \int_0^\infty \|\widehat{f}\chi_{E_{j,t}}\|_{L^2} \|\widehat{g}\chi_{E_{j,t}}\|_{L^2} \frac{dt}{t} \\ \leq & C\|\sigma_1\|_{L^2}^{4/5} 2^{-j\delta/5} \left(\int_0^\infty \int_{\mathbb{R}^n} |\widehat{f}\chi_{E_{j,t}}|^2 d\xi \frac{dt}{t} \right)^{\frac{1}{2}} \left(\int_0^\infty \int_{\mathbb{R}^n} |\widehat{g}\chi_{E_{j,t}}|^2 d\xi \frac{dt}{t} \right)^{\frac{1}{2}}.\end{aligned}$$

We control the last term as follows:

$$\int_0^\infty \int_{\mathbb{R}^n} |\widehat{f}\chi_{E_{j,t}}|^2 d\xi \frac{dt}{t} \leq C \int_{\mathbb{R}^n} \int_{1/|\xi|}^{2^j/|\xi|} \frac{dt}{t} |\widehat{f}(\xi)|^2 d\xi \leq Cj\|f\|_{L^2}^2$$

and thus we deduce

$$\|T(f,g)(x)\|_{L^1} \leq C\|\sigma_1\|_{L^2}^{4/5} 2^{-j\delta/5} j \|f\|_{L^2} \|g\|_{L^2}.$$

This completes the proof of Lemma 5. □

4. Proof of Proposition 4

Proof. Estimate (4) is automatically holds for finitely many terms in view of Proposition 3, so we fix a large j and define

$$T_{j,t}(f,g)(x) = \int_{\mathbb{R}^{2n}} \widehat{f}(\xi)\widehat{g}(\eta) m_j(t\xi, t\eta) e^{2\pi i x\cdot(\xi+\eta)} d\xi d\eta.$$

Take a smooth function ρ on $\mathbb{R}$ such that $\chi_{[\varepsilon-1,1-\varepsilon]} \leq \rho \leq \chi_{[-1,1]}$. Define $m_j^1(\xi,\eta) = m_j(\xi,\eta)\rho(\frac{1}{j}(\log_2 \frac{|\xi|}{|\eta|}))$, then we have a smooth decomposition of m_j with $m_j = m_j^1 + m_j^2$. On the support of m_j^1 we have $C^{-1}2^{-j}|\xi| \leq |\eta| \leq C2^j|\xi|$ and on the support of m_j^2 we have $2^{j(1-\varepsilon)}|\xi| \lesssim |\eta|$ or $2^{j(1-\varepsilon)}|\eta| \lesssim |\xi|$. We define

$$\mathcal{M}_j^i(f,g) = \sup_{t>0} |T_{j,t}^i(f,g)|, \quad i \in \{1,2\},$$

where $T_{j,t}^1$ and $T_{j,t}^2$ correspond to multipliers $m_j^1(t(\xi,\eta))$ and $m_j^2(t(\xi,\eta))$ respectively, such that $T_{j,t} = T_{j,t}^1 + T_{j,t}^2$. Then for f, g Schwartz functions we have

$$\mathcal{M}_j^1(f,g)(x) = \sup_{t>0} |T_{j,t}^1(f,g)(x)|$$

$$
\begin{aligned}
&= \sup_{t>0} \left| \int_0^t s \frac{dT^1_{j,s}(f,g)}{ds} \frac{ds}{s} \right| \\
&\le \int_0^\infty |\widetilde{T}^1_{j,s}(f,g)(x)| \frac{ds}{s},
\end{aligned}
$$

where $\widetilde{T}^1_{j,s}$ has bilinear multiplier $\widetilde{m}^1_j(s\xi,s\eta) = (s\xi,s\eta)\cdot(\nabla m^1_j)(s\xi,s\eta)$, a diagonal multiplier with nice decay, which can be used to establish the boundedness of the diagonal part with the aid of Lemma 5.

Recall that

$$
m^1_j(\xi,\eta) = \varphi(2^{-j}(\xi,\eta)) 2\pi \frac{J_{n-1}(2\pi(\xi,\eta))}{|(\xi,\eta)|^{n-1}} \rho(\tfrac{1}{j}(\log_2 \tfrac{|\xi|}{|\eta|}))
$$

for $j \ge 1$ and a calculation shows that $|\partial_1(m^1_j)|$ is controlled by the sum of three terms bounded by $C2^{-j(2n-1)/2}$, $C2^{-j(2n+1)/2}$ and $C\frac{1}{j}2^{-j(2n-1)/2}$ respectively. Indeed, when the derivative falls on ϕ, we can bound it by $C2^{-j}2^{-j(n-1/2)} = C2^{-j(n+1/2)}$. If the derivative falls on the second part, using properties of Bessel functions (see, e.g., [9, Appendix B.2]), we obtain the bound $C\frac{J_n(2\pi(\xi,\eta))}{|(\xi,\eta)|^n}|\xi_1| \le C2^{-j(n-1/2)}$. For the last case, we can bound it by $C2^{-j(n-1/2)} j^{-1} \frac{1}{|\xi|}\frac{\xi_1}{|\xi|} \le C2^{-j(n-1/2)} j^{-1} 2^{-\varepsilon j}$. As a consequence we have $|\partial_1(m^1_j)| \le C2^{-j(2n-1)/2}$. Then we can show that $|\partial_1(\widetilde{m}^1_j)| \le C2^{-j(2n-3)/2}$ and similar arguments give that for any multiindex α we have $|\partial^\alpha \widetilde{m}^1_j| \le C2^{-j(2n-3)/2}$. Moreover, from this we can show that

$$
\|\widetilde{m}^1_j\|_2 \le C\left(\int_{|(\xi,\eta)|\sim 2^j} |2^{-j(n-\frac{3}{2})}|^2 d\xi d\eta \right)^{\frac{1}{2}} \le C2^{-j(n-\frac{3}{2})} 2^{jn} \le C2^{\frac{3}{2}j}.
$$

Applying Lemma 5 to the function $\widetilde{m}^1_j(\xi,\eta) = (\xi,\eta)\cdot(\nabla m^1_j)(\xi,\eta)$ which satisfies the hypotheses with $\delta = (2n-3)/2$, we obtain

$$
\|\mathcal{M}^1_j(f,g)\|_{L^1} \le Cj \|\widetilde{m}^1_j\|_{L^2}^{\frac{4}{5}} 2^{-j\frac{\delta}{5}} \|f\|_{L^2}\|g\|_{L^2} = Cj2^{j(\frac{3}{2}-\frac{n}{5})}\|f\|_{L^2}\|g\|_{L^2}. \tag{5}
$$

It remains to obtain an analogous estimate for $\mathcal{M}^2_j$.

For the off-diagonal part m^2_j we use a different decomposition involving g-functions. For $f,g \in \mathcal{S}(\mathbb{R}^n)$ we have

$$
\begin{aligned}
\mathcal{M}^2_j(f,g)(x) &= \big(\sup_{t>0} |T^2_{j,t}(f,g)(x)|^2\big)^{\frac{1}{2}} \\
&= \left(\sup_{t>0} \left| 2\int_0^t T^2_{j,s}(f,g)(x)\, s \frac{dT^2_{j,s}(f,g)(x)}{ds} \frac{ds}{s} \right| \right)^{\frac{1}{2}} \\
&\le \sqrt{2} \Big\{ \Big(\int_0^\infty |T^2_{j,s}(f,g)|^2 \frac{ds}{s} \Big)^{\frac{1}{2}} \Big(\int_0^\infty |\widetilde{T}^2_{j,s}(f,g)|^2 \frac{ds}{s} \Big)^{\frac{1}{2}} \Big\}^{\frac{1}{2}}
\end{aligned}
$$

$$= \sqrt{2}\left(G_j(f,g)(x)\widetilde{G}_j(f,g)\right)^{\frac{1}{2}}. \tag{6}$$

Here $\widetilde{T}^2_{j,s}(f,g)$ has symbol $\widetilde{m}^2_j(s\xi,s\eta) = (s\xi,s\eta)\cdot(\nabla m^2_j)(s\xi,s\eta)$ and

$$\begin{aligned} G_j(f,g)(x) &= \Big(\int_0^\infty |T^2_{j,s}(f,g)|^2 \frac{ds}{s}\Big)^{\frac{1}{2}} \\ \widetilde{G}_j(f,g)(x) &= \Big(\int_0^\infty |\widetilde{T}^2_{j,s}(f,g)|^2 \frac{ds}{s}\Big)^{\frac{1}{2}}. \end{aligned}$$

Lemma 6. *If a $\sigma_1(\xi,\eta)$ on $\mathbb{R}^{2n}$ satisfies*

(i) for any multiindex $|\alpha| \le M = 4n$, there exists a positive constant C_α independent of j such that $\|\partial^\alpha(\sigma_1(\xi,\eta))\|_{L^\infty} \le C_\alpha 2^{-j\delta}$,

(ii) supp $\sigma_1 \subset \{(\xi,\eta)\in\mathbb{R}^{2n} : |(\xi,\eta)| \sim 2^j, |\xi| \ge 2^{j(1-\varepsilon)}|\eta|$, or $|\eta| \ge 2^{j(1-\varepsilon)}|\xi|\}$,

then $T(f,g)(x) := (\int_0^\infty |T_{\sigma_t}(f,g)(x)|^2 \frac{dt}{t})^{1/2}$ is bounded from $L^2\times L^2$ to L^1 with bound at most a multiple of $2^{-j(\delta-\varepsilon)}$, where $\sigma_t(\xi,\eta) = \sigma_1(t\xi,t\eta)$.

Proof. Recall that supp $m^2_j \subset \{(\xi,\eta) : 2^{j(1-\varepsilon)}|\xi| \lesssim |\eta|$ or $2^{j(1-\varepsilon)}|\eta| \lesssim |\xi|\}$. We consider only the part $\{|\xi| \ge 2^{j(1-\varepsilon)}|\eta|\}$ because the other part is similar. By [10, Section 5] we have

$$|T_{\sigma_1}(f,g)(x)| \le C2^{\varepsilon j}2^{-j\delta}M(g)(x)|T_m(f)(x)|,$$

where M is the Hardy-Littlewood maximal function and T_m is a linear operator that satisfies $\|T_m(f)\|_{L^2} \le C\|\widehat{f}\chi_{\{|\xi|\sim 2^j\}}\|_{L^2}$. Then

$$|T_{\sigma_t}(f,g)(x)| \le 2^{-j(\delta-\varepsilon)}t^{-n}M(g)(x)T_m(f_t)(x/t),$$

and

$$\begin{aligned} &\int_{\mathbb{R}^n}\left(\int_0^\infty |T_{\sigma_t}(f,g)(x)|^2\frac{dt}{t}\right)^{\frac{1}{2}}dx \\ &\quad\le C2^{-j(\delta-\varepsilon)}\int_{\mathbb{R}^n}\left(\int_0^\infty t^{-2n}M(g)(x)^2|T_m(f_t)(x/t)|^2\frac{dt}{t}\right)^{\frac{1}{2}}dx \\ &\quad\le C2^{-j(\delta-\varepsilon)}\|M(g)\|_{L^2}\Big(\int_{\mathbb{R}^n}\int_0^\infty \big|t^{-n}T_m(f_t)(x/t)\big|^2\frac{dt}{t}dx\Big)^{\frac{1}{2}} \\ &\quad\le C2^{-j(\delta-\varepsilon)}\|g\|_{L^2}\Big(\int_{\mathbb{R}^n}|\widehat{f}(\xi)|^2\int_{2^{j-1}/|\xi|}^{2^{j+1}/|\xi|}\frac{dt}{t}d\xi\Big)^{\frac{1}{2}} \\ &\quad\le C2^{-j(\delta-\varepsilon)}\|g\|_{L^2}\|f\|_{L^2}. \end{aligned}$$

This completes the proof of Lemma 6. □

We now return to the proof of Proposition 4. Notice that both $m_j^2(\xi,\eta)$ and $\widetilde{m}_j^2(\xi,\eta)$ satisfy conditions of Lemma 6 with δ being either $(2n-1)/2$ or $(2n-3)/2$ respectively, so

$$\|G_j(f,g)\|_{L^1} \le C2^{-j(2n-1)/2}\|f\|_{L^2}\|g\|_{L^2}$$
$$\|\widetilde{G}_j(f,g)\|_{L^1} \le C2^{-j(2n-3)/2}\|f\|_{L^2}\|g\|_{L^2}.$$

Using (6) we deduce

$$\|\mathcal{M}_j^2(f,g)\|_{L^1} \le \|G_j(f,g)\|_{L^1}^{1/2}\|\widetilde{G}_j(f,g)\|_{L^1}^{1/2} \le C2^{-j(n-1)}\|f\|_{L^2}\|g\|_{L^2}. \tag{7}$$

Combining (5) and (7) yields Proposition 4 with $\delta_n = \frac{n}{5} - \frac{3}{2}$. □

5. INTERPOLATION

By Proposition 3 (for term $j \le c_0$) and Proposition 4 (for $j \ge c_0$), for any $\delta_n' < \delta_n$, as a consequence of (3) we obtain

$$\|\mathcal{M}(f,g)\|_{L^1} \le \sum_{j=0}^{\infty} C_{\delta_0'}2^{-\delta_n' j}\|f\|_{L^2}\|g\|_{L^2} \le C_{\delta_0'}\|f\|_{L^2}\|g\|_{L^2}.$$

This establishes the boundedness of $\mathcal{M}$ from $L^2 \times L^2$ to L^1 claimed in Theorem 1 (recall $n \ge 8$). It remains to obtain estimates for other values of p_1, p_2. This is achieved via bilinear interpolation.

Notice that when one index among p_1 and p_2 is equal to 1, we have that $\mathcal{M}_j$ maps $L^{p_1} \times L^{p_2}$ to $L^{p,\infty}$ with norm $\lesssim 2^j$. Indeed, this follows from the estimate

$$|\varphi_j^\vee * (d\sigma)(y,z)| \le C_N 2^j(1+|(y,z)|)^{-2N} \le C_N 2^j(1+|y|)^{-N}(1+|z|)^{-N}$$

which can be found, for instance, in [9, estimate (6.5.12)]. Thus we have

$$\mathcal{M}_j(f,g)(x) \le C2^j M(f)M(g)$$

where M is the Hardy-Littlewood maximal function. We pick two points

$$\vec{Q}_1 = (1/1, 1/(1+\varepsilon), (2+\varepsilon)/(1+\varepsilon))$$
$$\vec{Q}_2 = (1/(1+\varepsilon), 1/1, (2+\varepsilon)/(1+\varepsilon))$$

and we also consider the point $\vec{Q}_0 = (1/2, 1/2, 1)$. We interpolate the known estimates for $\mathcal{M}_j$ at these three points. Letting ε go to 0, we obtain that for $p > \frac{2+2\delta_n}{1+2\delta_n}$ we have that $\mathcal{M}_j$ maps $L^p(\mathbb{R}^n) \times L^p(\mathbb{R}^n)$ to $L^{p/2}(\mathbb{R}^n)$ with a geometrically decreasing bound in j. Recall that $\delta_n = (2n-15)/10 > 0$, so we need $n \ge 8$.

Thus summing over j gives boundedness for $\mathcal{M}$ from $L^p(\mathbb{R}^n) \times L^p(\mathbb{R}^n)$ to $L^{p/2}(\mathbb{R}^n)$ when $p > \frac{2+2\delta_n}{1+2\delta_n}$. By interpolation we obtain boundedness for $\mathcal{M}$ in the interior of a rhombus with vertices the points $(1/\infty, 1/\infty, 1/\infty)$,

$(\frac{2n-3/2}{2n-1}, \frac{1}{\infty}, \frac{2n-3/2}{2n-1})$, $(\frac{1}{\infty}, \frac{2n-3/2}{2n-1}, \frac{2n-3/2}{2n-1})$ and $(\frac{1+2\delta_n}{2+2\delta_n}, \frac{1+2\delta_n}{2+2\delta_n}, \frac{2+4\delta_n}{2+2\delta_n})$. The proof of Theorem 1 is now complete.

We remark that is the largest region for which we presently know boundedness for $\mathcal{M}$ in dimensions $n \geq 8$.

6. COUNTEREXMAPLES

In this section we construct counterexamples indicating the unboundedness of the bilinear spherical maximal operator in a certain range. Our examples are inspired by Stein [19] but the situation is more complicated.

Proposition 7. *The bilinear spherical maximal operator $\mathcal{M}$ is unbounded from $L^{p_1}(\mathbb{R}^n) \times L^{p_2}(\mathbb{R}^n)$ to $L^p(\mathbb{R}^n)$ when $1 \leq p_1, p_2 \leq \infty$, $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$, $n \geq 1$, and $p \leq \frac{n}{2n-1}$. In particular, $\mathcal{M}$ is unbounded from $L^2(\mathbb{R}) \times L^2(\mathbb{R})$ to $L^1(\mathbb{R})$ when $n = 1$.*

Remark 1. We note that $\frac{1+\delta_n}{1+2\delta_n} - \frac{n}{2n-1} = \frac{1+\frac{n}{5}-\frac{3}{2}}{1+\frac{2n}{5}-3} - \frac{n}{2n-1} \approx \frac{1}{n} \to 0$ as $n \to \infty$. This means that the gap between the range of boundedness and unboundedness tends to 0 as the dimension increases to infinity.

Proof. We first consider the case $n = 1$ where it is easy to demonstrate the main idea.

Define functions on $\mathbb{R}$ by setting $f(y) = |y|^{-1/p_1}(\log \frac{1}{|y|})^{-2/p_1} \chi_{|y|\leq 1/2}$ and $g(y) = |y|^{-1/p_2}(\log \frac{1}{|y|})^{-2/p_2} \chi_{|y|\leq 1/2}$. Then $f \in L^{p_1}(\mathbb{R})$, $g \in L^{p_2}(\mathbb{R})$ and we will estimate from below $M_{\sqrt{2}R}(f,g)(R)$ for large R, where

$$M_t(f,g)(x) = \int_{\mathbb{S}^1} |f(x-ty)g(x-tz)| d\sigma(y,z).$$

In view o the support properties of f and g we have $|y - \frac{1}{\sqrt{2}}| \leq \frac{1}{2\sqrt{2}R}$, and $|z - \frac{1}{\sqrt{2}}| \leq \frac{1}{2\sqrt{2}R}$. We also have that $y^2 + z^2 = 1$ since $(y,z) \in \mathbb{S}^1$.

Therefore we rewrite $M_{\sqrt{2}R}(f,g)(R)$ as

$$\int_{\frac{\sqrt{2}}{2}-\frac{1}{2\sqrt{2}R}}^{\frac{\sqrt{2}}{2}+\frac{1}{2\sqrt{2}R}} |R(1-\sqrt{2}y)|^{-\frac{1}{p_1}} (-\log|R(1-\sqrt{2}y)|)^{-\frac{2}{p_1}} \tag{8}$$
$$|R(1-\sqrt{2}z)|^{-\frac{1}{p_2}} (-\log|R(1-\sqrt{2}z)|)^{-\frac{2}{p_2}} \frac{dy}{\sqrt{1-y^2}},$$

with $z = \sqrt{1-y^2}$.

Notice that $|R(1-\sqrt{2}z)| = R|\frac{1-2z^2}{1+\sqrt{2}z}| \leq R|1-2y^2| \leq 3R|1-\sqrt{2}y|$ since[1] $z \approx y \approx \sqrt{2}/2$. As a result, with the help of (9) [Lemma 8], the expression

[1]Here $a \approx b$ means that $|a-b|$ is very small.

in (8) is greater than

$$\int_{\frac{\sqrt{2}}{2}-\frac{1}{100R}}^{\frac{\sqrt{2}}{2}+\frac{1}{100R}} R^{-\frac{1}{p}}|(1-\sqrt{2}y)|^{-\frac{1}{p}}(-\log|R(1-\sqrt{2}y)|)^{-\frac{2}{p}}dy$$
$$=2R^{-1}\int_0^{\frac{1}{100}} t^{-1/p}(\log\tfrac{1}{t})^{-2/p}dt = \begin{cases} C_pR^{-1} & \text{if } p\geq 1 \\ \infty & \text{if } p<1. \end{cases}$$

Thus $\mathcal{M}(f,g)\notin L^p(\mathbb{R})$ for $p<1$ and also $\mathcal{M}(f,g)(x)\geq C/x$ for x large if $p=1$. It follows that $\mathcal{M}(f,g)\notin L^1(\mathbb{R})$ for $p=1$, hence the statement of the proposition holds.

We now consider the higher-dimensional case $n\geq 2$. We define $f(y)=|y|^{-n/p_1}(\log\frac{1}{|y|})^{-2/p_1}\chi_{|y|\leq 1/100}$ and $g(y)=|y|^{-n/p_2}(\log\frac{1}{|y|})^{-2/p_2}\chi_{|y|\leq 1/2}$. We have that f lies in $L^{p_1}(\mathbb{R}^n)$ and g lies in $L^{p_2}(\mathbb{R}^n)$. The mapping $(y,z)\mapsto(Ay,Az)$ with $A\in SO_n$ is an isometry on $\mathbb{S}^{2n-1}$, hence we have $M_t(f,g)(x)=M_t(f,g)(|x|e_1)$, where $e_1=(1,0,\dots,0)\in\mathbb{R}^n$. Thus we may take $x=Re_1\in\mathbb{R}^n$ with R large.

By the change of variables identity (10) [Lemma 9], we have

$$\begin{aligned} &M_{\sqrt{2}R}(f,g)(Re_1)\\ &\quad=\int_{\mathbb{S}^{2n-1}} f(Re_1-\sqrt{2}Ry)g(Re_1-\sqrt{2}Rz)d\sigma(y,z)\\ &\quad=\int_{B_n(\frac{1}{\sqrt{2}}e_1,\frac{1}{100R})}|\sqrt{R}y-Re_1|^{-\frac{n}{p_1}}(-\log|Re_1-\sqrt{2}Ry|)^{-\frac{2}{p_1}}\\ &\qquad\int_E|\sqrt{2}Rz-Re_1|^{-\frac{n}{p_2}}(-\log|Re_1-\sqrt{2}Rz|)^{-\frac{2}{p_2}}d\sigma_{n-1}^r(z)\tfrac{dy}{\sqrt{1-|y|^2}}, \end{aligned}$$

where $B_n(a,r)$ is a ball in $\mathbb{R}^n$ centered at a with radius r, and E is the $(n-1)$-dimensional manifold $\mathbb{S}^{n-1}_{\sqrt{1-|y|^2}}\cap B_n(\frac{1}{\sqrt{2}}e_1,\frac{1}{2\sqrt{2}R})$ with $\mathbb{S}^{n-1}_r$ being the sphere in $\mathbb{R}^n$ with radius r and $d\sigma_{n-1}^r$ the measure on $\mathbb{S}^{n-1}_r$.

We next focus on the inner integral, namely

$$I=\int_E|\sqrt{2}Rz-Re_1|^{-\frac{n}{p_2}}(-\log|Re_1-\sqrt{2}Rz|)^{-\frac{2}{p_2}}d\sigma_{n-1}^r(z).$$

Take a point $z_0\in\mathbb{S}^{n-1}_{\sqrt{1-|y|^2}}\cap\partial\big(B_n(\frac{1}{\sqrt{2}}e_1,\frac{1}{2\sqrt{2}R})\big)$, and let θ be the angle between vectors z_0 and e_1, which the largest one between $z\in E$ and e_1. Here ∂B is the boundary of a set B. Then θ is small if R is large and[2] $|E|\sim(\sqrt{1-|y|^2}\theta)^{n-1}\sim\theta^{n-1}$. Noticing that $\theta^2\sim\sin^2\theta=1-\cos^2\theta\sim$

[2] $A\sim B$ means that the ratio A/B is bounded above and below

$1-\cos\theta$ and that

$$1-|y|^2+\tfrac{1}{2}-\sqrt{2}\sqrt{1-|y|^2}\cos\theta=\tfrac{1}{8R^2},$$

we obtain that $\theta^2\sim\frac{1}{8R^2}-(\sqrt{1-|y|^2}-\frac{1}{\sqrt{2}})^2$. Then we write

$$\Big|\sqrt{1-|y|^2}-\tfrac{1}{\sqrt{2}}\Big|=\Big|\tfrac{1-|y|^2-\frac{1}{2}}{\sqrt{1-|y|^2}+\frac{1}{\sqrt{2}}}\Big|\leq 2|\tfrac{1}{2}-|y|^2|\leq\tfrac{1}{25R}.$$

Consequently $\theta\geq C/R$.

Collecting the previous calculations, we can bound I from below by

$$\int_0^\theta\int_{\mathbb{S}^{n-2}_{t\sin\alpha}}|\sqrt{2}Rz-Re_1|^{-\frac{n}{p_2}}(-\log|Re_1-\sqrt{2}Rz|)^{-\frac{2}{p_2}}d\sigma^{t\sin\alpha}_{n-2}(z)d\alpha,$$

where $t=|z|=\sqrt{1-|y|^2}\approx\frac{1}{\sqrt{2}}$, and $z_1=\cos\alpha$. By symmetry, let us consider just that case $t<\frac{1}{\sqrt{2}}$. Let β be the angle such that $|\sqrt{2}z-e_1|=2|\sqrt{2}t-1|$, then $2t^2+1-2\sqrt{2}t\cos\beta=4|\sqrt{2}t-1|^2$, which implies that $\beta^2\sim 1-\cos\beta\sim 2\sqrt{2}t-2t^2-1+4(\sqrt{2}t-1)^2=3(\sqrt{2}t-1)^2$. So $\beta\sim 1-\sqrt{2}t$. When $\alpha=0$, we have trivially that $|\sqrt{2}z-e_1|=|\sqrt{2}t-1|$. So for $\alpha\in[0,\beta]$, we have $|\sqrt{2}z-e_1|\sim 2|\sqrt{2}t-1|\leq 2\big|2|z|^2-1\big|=2\big|2|y|^2-1\big|\leq 6\big|\sqrt{2}|y|-1\big|\leq 6|\sqrt{2}y-e_1|$. Consequently using the fact that $1-\sqrt{2}t\leq C\theta$ and (9) again we obtain

$$\begin{aligned}
I&\geq C\int_0^\theta\int_{\mathbb{S}^{n-2}_{t\sin\alpha}}\frac{|\sqrt{2}Rz-Re_1|^{1-n}}{|\sqrt{2}Rz-Re_1|^{\frac{n}{p_2}-n+1}(-\log|Re_1-\sqrt{2}Rz|)^{\frac{2}{p_2}}}\,d\sigma^{t\sin\alpha}_{n-2}(z)\,d\alpha\\
&\geq\frac{CR^{1-n}|\sqrt{2}t-1|^{1-n}}{|\sqrt{2}Ry-Re_1|^{\frac{n}{p_2}-n+1}(-\log|Re_1-\sqrt{2}Ry|)^{\frac{2}{p_2}}}\int_0^{C(1-\sqrt{2}t)}\sin^{n-2}\alpha d\alpha\\
&\geq CR^{1-n}\frac{|\sqrt{2}t-1|^{1-n}|1-\sqrt{2}t|^{n-1}}{|\sqrt{2}Ry-Re_1|^{\frac{n}{p_2}-n+1}(-\log|Re_1-\sqrt{2}Ry|)^{\frac{2}{p_2}}}\\
&=CR^{1-n}|\sqrt{2}Ry-Re_1|^{-\frac{n}{p_2}+n-1}(-\log|Re_1-\sqrt{2}Ry|)^{-\frac{2}{p_2}}.
\end{aligned}$$

Using this estimate we see that

$$\begin{aligned}
&M_{\sqrt{2}R}(f,g)(Re_1)\\
&\quad\geq CR^{1-n}\int_{B_n(\frac{1}{\sqrt{2}}e_1,\frac{1}{100R})}|Re_1-\sqrt{2}Ry|^{-\frac{n}{p}+n-1}(-\log|Re_1-\sqrt{2}Ry|)^{-\frac{2}{p}}dy\\
&\quad=CR^{1-2n}\int_{B_n(0,\frac{1}{100})}|x|^{-\frac{n}{p}+n-1}(-\log|x|)^{-\frac{2}{p}}dx
\end{aligned}$$

$$= CR^{1-2n}\int_0^{\frac{1}{100}} r^{-\frac{n}{p}+2n-2}(-\log r)^{-\frac{2}{p}}dr$$
$$= \begin{cases} CR^{-2n+1} & \text{if } p = \frac{n}{2n-1} \\ \infty & \text{if } p < \frac{n}{2n-1} \end{cases}.$$

Hence $\mathcal{M}(f,g)$ is not in L^p for $p < \frac{n}{2n-1}$ and $\mathcal{M}(f,g)(x) \geq C|x|^{1-2n}$ for all $|x|$ large enough, hence it is also not in $L^{\frac{n}{2n-1}}(\mathbb{R}^n)$ when $p = \frac{n}{2n-1}$. □

Lastly, we prove a couple of points left open.

Lemma 8. *Let $r_1, r_2 > 0$, $t,\ s \leq \frac{1}{10}$, and $t \leq Cs$ for some $C \geq 1$. Then there exists an absolute constant C' (depending on C, r_1, r_2) such that*

$$s^{-r_1}(\log\tfrac{1}{s})^{-r_2} \leq C' t^{-r_1}(\log\tfrac{1}{t})^{-r_2}. \tag{9}$$

Proof. Define $F(x) = x^{r_1}(\log x)^{-r_2}$. Differentiating F, we see that F is increasing when x is large enough and so,

$$F(\tfrac{1}{s}) = s^{-r_1}(\log\tfrac{1}{s})^{-r_2} \leq C^{r_1}(Cs)^{-r_1}(\log\tfrac{1}{Cs})^{-r_2} = C^{r_1}F(\tfrac{1}{Cs}) \leq C'F(\tfrac{1}{t}),$$

which is a restatement of (9).

□

Lemma 9. *For functions $F(y,z)$ defined in $\mathbb{R}^{2n}$ with $y,\ z \in \mathbb{R}^n$, we have*

$$\int_{\mathbb{S}^{2n-1}} F(y,z)d\sigma(y,z) = \int_{B_n}\int_{\mathbb{S}^{n-1}_{r_y}} F(y,z)d\sigma^{r_y}_{n-1}(z)\frac{dy}{\sqrt{1-|y|^2}}, \tag{10}$$

where B_n is the unit ball in $\mathbb{R}^n$ and $\mathbb{S}^{n-1}_{r_y}$ is the sphere in $\mathbb{R}^n$ centered at 0 with radius $r_y = \sqrt{1-|y|^2}$.

Proof. We begin by writing $\int_{\mathbb{S}^{2n-1}} F(y,z)d\sigma(y,z)$ as

$$\int_{B_{2n-1}} \big[F(y,z',z_n) + F(y,z',-z_n)\big]\frac{dydz'}{\sqrt{1-|y|^2-|z'|^2}}, \tag{11}$$

where $z = (z', z_n)$, and $z_n = \sqrt{1-|y|^2-|z'|^2}$; see [9, Appendix D.5].

Writing $z/r_y = \omega = (\omega', \omega_{n-1}) \in \mathbb{R}^{n-1}\times\mathbb{R}$, we express the right hand side of (10) as

$$\int_{B_n}\int_{\mathbb{S}^{n-1}_{r_y}} F(y,z)d\sigma^{r_y}_{n-1}(z)\frac{dy}{\sqrt{1-|y|^2}}$$
$$= \int_{B_n} r_y^{n-1}\int_{\mathbb{S}^{n-1}} F(y,r_y\omega)d\sigma_{n-1}(\omega)\frac{dy}{\sqrt{1-|y|^2}}$$
$$= \int_{B_n} r_y^{n-1}\int_{B_{n-1}} \big[F(y,r_y\omega',r_y\omega_n) + F(y,r_y\omega',-r_y\omega_n)\big]\frac{d\omega'}{\sqrt{1-|\omega'|^2}}\frac{dy}{\sqrt{1-|y|^2}}$$

$$= \int_{B_n} r_y^{n-1} \int_{r_y B_{n-1}} \big[F(y,z',z_n) + F(y,z',-z_n)\big] \frac{r_y^{1-n} dz'}{\sqrt{1-|\omega'|^2}} \frac{dy}{\sqrt{1-|y|^2}}$$

$$= \int_{B_n} \int_{r_y B_{n-1}} \big[F(y,z',z_n) + F(y,z',-z_n)\big] \frac{dy dz'}{\sqrt{1-|y|^2-|z'|^2}},$$

as one can easily verify that $\sqrt{1-|\omega'|^2}\sqrt{1-|y|^2} = \sqrt{1-|y|^2-|z'|^2}$. Using that B_{2n-1} is equal to the disjoint union of the sets $\{(y, r_y v) : v \in B_{n-1}\}$ over all $y \in B_n$, we see that the last double integral is equal to the expression in (11), as claimed. □

REFERENCES

[1] Bourgain, J., *Averages in the plane over convex curves and maximal operators*. Journal d'Analyse Mathématique **47**(1) (1986), 69–85.

[2] Calderón, C. P., *Lacunary spherical means*. Illinois Journal of Mathematics **23**(3) (1979), 476–484.

[3] Carbery, A., *Radial Fourier multipliers and associated maximal functions*. North-Holland Mathematics Studies **111** (1985), 49–56.

[4] Coifman, R. R. and Weiss, G., Review: R. E. Edwards and G. Gaudry, *Littlewood-Paley and multiplier theory*. Bulletin of the American Mathematical Society **84**(2) (1978), 242–250.

[5] Cowling, M. and Mauceri, G., *On maximal functions*. Milan Journal of Mathematics **49**(1) (1979), 79–87.

[6] Duoandikoetxea, J. and Rubio de Francia, J.-L., *Maximal and singular integral operators via Fourier transform estimates*. Inventiones Mathematicae **84** (1986), 541–561.

[7] Duoandikoetxea, J. and Vega, L., *Spherical means and weighted inequalities*. Journal of the London Mathematical Society **53**(2) (1996), 343–353.

[8] Geba, D., Greenleaf, A., Iosevich, A., Palsson, E. and Sawyer, E. *Restricted convolution inequalities, multilinear operators and applications*. Mathematical Research Letters **20** (2013), no. 4, 675–694.

[9] Grafakos, L., *Classical Fourier Analysis*, Third Edition. Graduate Texts in Mathematics, **249**, Springer, New York, 2014.

[10] Grafakos, L., He, D., and Honzík P., *Rough bilinear singular integrals*. Submitted (2016).

[11] Grafakos, L., He, D., and Honzík P., *The Hörmander multiplier theorem, II: The bilinear local L^2 case*. Submitted (2016).

[12] Greenleaf, A., *Principal curvature and harmonic-analysis*. Indiana University Mathematics Journal **30**(4) (1981), 519–537.

[13] Greenleaf, A., Iosevich, A., Krause, B., and Liu, A., *Bilinear generalized Radon transforms in the plane*. https://arxiv.org/abs/1704.00861

[14] He, D., *On bilinear maximal Bochner-Riesz operators*. Submitted (2016).

[15] Magyar, A., Stein, E., and Wainger, S., *Discrete analogues in harmonic analysis: spherical averages*. Annals of Mathematics (2nd Ser.) **155**(1) (2002), 189–208.

[16] Mockenhaupt, G., Seeger, A., and Sogge, C. D., *Wave front sets, local smoothing and Bourgain's circular maximal theorem*. Annals of Mathematics (2nd Ser.) **136**(1) (1992), 207–218.

[17] Rubio de Francia, J. L., *Maximal functions and Fourier transforms*. Duke Mathematical Journal **53**(2) (1986), 395–404.
[18] Schlag, W., *A geometric proof of the circular maximal theorem*. Duke Mathematical Journal **93**(3) (1998), 505–534.
[19] Stein, E. M., *Maximal functions: spherical means*. Proceedings of the National Academy of Sciences, **73**(7) (1976), 2174–2175.
[20] Stein, E. M., *Harmonic Analysis, Real Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton Mathematical Series, 43, Monographs in Harmonic Analysis, III, Princeton University Press, Princeton, NJ, 1993.

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL PORTO ALEGRE, RS, BRAZIL 91509-900
E-mail address: josea@mat.ufrgs.br

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211, USA
E-mail address: grafakosl@missouri.edu

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, P. R. CHINA
E-mail address: hedanqing@mail.sysu.edu.cn

MFF UK, SOKOLOVSKA 83, PRAHA 7, CZECH REPUBLIC
E-mail address: honzik@gmail.com

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL PORTO ALEGRE, RS, BRAZIL 91509-900
E-mail address: lucas.oliveira@ufrgs.br